

## Antibody Data

<b>Product SKU:</b>	<b>AGEL0046</b>	<b>Clone:</b>	<b>Cl:A3-1</b>
<b>Applications:</b>	<b>FCM</b>		
<b>Reactivity:</b>	<b>Mouse</b>		

## Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

## Product Information:

**Alternate Names:** Adhesion G protein-coupled receptor E1;Adgre1;Cell surface glycoprotein F4/80;EGF-like module receptor 1;Adgre1;Emr1; Gpf480;

**Uniprot ID:** Q61549

**Background:** F4/80 is a 160 kD glycoprotein. It is characterized as a member of the epidermal growth factor (EGF)-transmembrane 7 (TM7) family. F4/80, also known as EMR1 or Ly71, has been widely used as a murine macrophage marker, which is expressed on the majority of tissue macrophages including peritoneal macrophages, macrophages in lung, gut, thymus and red pulp of spleen (but not on the macrophages located in T cell areas of the spleen, lymph node and Peyer's patch), Kuffer cells, Langerhans cells, and bone marrow stromal cells. F4/80 has also been shown on a subset of dendritic cells. The biological ligand of F4/80 has not been identified, but it has been reported that F4/80 is required for induction of CD8+ T cells-mediated peripheral tolerance.


**Form:** Liquid

**Conjugation:** APC

**Size:** 50 Tests, 100 Tests, 200 Tests

**Host Species:** Rat

**Isotype:** Rat IgG2b,  $\kappa$



Ex:650 nm; Em:660 nm

**Isotype Control:** APC Rat IgG2b,  $\kappa$  Isotype Control[LTF-2] [Product AGEL0046]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C

**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

**Recommended Usage:** Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

---