

**Biotin Anti-Mouse CD54 Antibody  
[YN1/1.7.4]**

Catalogue Code: AGEL0117

**Antibody Data**

<b>Product SKU:</b>	<b>AGEL0117</b>	<b>Clone:</b>	<b>YN1/1.7.4</b>
<b>Applications:</b>	<b>FCM</b>		
<b>Reactivity:</b>	<b>Mouse</b>		

**Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

**Product Information:**

<b>Alternate Names:</b>	Intercellular adhesion molecule 1;Icam1;MALA-2;MyD10;CD54;Icam-1;
<b>Uniprot ID:</b>	P13597
<b>Background:</b>	CD54 is a 90 kD immunoglobulin superfamily member also known as ICAM-1 and Ly-47. It is expressed on activated endothelial cells, high endothelial venules (HEV), T and B cells, monocytes/ macrophages, granulocytes, and dendritic cells. CD54 is an important intracellular adhesion molecule that participates in T cell-T cell, T cell-B cell, and T cell-target cell interactions via binding of LFA-1 (CD11a/CD18) and Mac-1 (CD11b/CD18). CD54 has also been shown to be involved in lymphocyte trafficking, making it an important molecule in many immune reactions and inflammation. CD54 is also a receptor for rhinovirus. The YN1/1.7.4 antibody has been reported to block binding of mouse CD54 to LFA-1 and Mac-1, inhibit cell-cell adhesion, and function in antigen presentation to T cells and leukocyte migration to inflammatory tissues.
<b>Form:</b>	Liquid
<b>Conjugation:</b>	Biotin
<b>Size:</b>	25&micro;g, 100&micro;g
<b>Host Species:</b>	Rat
<b>Isotype:</b>	Rat IgG2b, $\kappa$
<b>Isotype Control:</b>	Biotin Rat IgG2b, $\kappa$ Isotype Control[LTF-2] [Product AGEL0117]
<b>Storage Buffer:</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
<b>Shipping:</b>	Biological ice pack at 4°C

**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

**Recommended Usage:** Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 1.0 \mu\text{g}$  per  $10^6$  cells in 100  $\mu\text{L}$  volume or 100  $\mu\text{L}$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

---