

**GenieFluor 488 Anti-Human CD41  
Antibody [HIP8]  
Catalogue Code: AGEL0327**

**Antibody Data**

<b>Product SKU:</b>	<b>AGEL0327</b>	<b>Clone:</b>	<b>HIP8</b>
<b>Applications:</b>	<b>FCM</b>		
<b>Reactivity:</b>	<b>Human</b>		

**Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

**Product Information:**

**Alternate Names:** Integrin alpha-IIb/3;ITGA2B;GPIIb/3;GPIIb;Platelet membrane glycoprotein IIb/3;CD41;GP2B; ITGAB;

**Uniprot ID:** P08514

**Background:** CD41 is a 125/25 kD  $\alpha$  subunit of the GPIIb/IIIa (CD41/CD61) complex. CD41 is a heterodimer composed of a heavy chain (GPIIb $\alpha$ ) and light chain (GPIIb $\beta$ ) linked by a single disulfide bond. It is a member of the integrin family primarily expressed on platelets and megakaryocytes. CD41 has been reported to be involved with platelet aggregation and platelet attachment to the ECM. CD41/CD61 complex acts as the receptor for fibrinogen, fibronectin, Von Willebrand factor and thrombin.


**Form:** Liquid

**Conjugation:** Genie Fluor488

**Size:** 20 Tests, 100 Tests, 200 Tests

**Host Species:** Mouse

**Isotype:** Mouse IgG1,  $\kappa$



**Isotype Control:** Genie Fluor 488 Mouse IgG1,  $\kappa$  Isotype Control[MOPC-21] [Product AGEL0327]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C

**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

**Recommended Usage:** Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.