

Product Datasheet

Purified Anti-Mouse CD103 Antibody [M290]

Catalogue Code: AGEL0330

Antibody Data

Product SKU: AGEL0330 Clone: M290

Applications: FCM

Reactivity: Mouse

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: Integrin alpha-E;Itgae;Integrin alpha M290;CD103;

Uniprot ID: Q60677

Background: CD103 is a type I transmembrane glycoprotein known as αE integrin or Integrin αIEL chain.

It belongs to the integrin family and is primarily found on intestinal intraepithelial lymphocytes (IEL). CD103 is also expressed on a subpopulation of lamina propria T cells, epithelial dendritic cells, lamina propria-derived dendritic cells, and a small subset of peripheral lymphocytes. T regulatory cells express high level of CD103. The CD103 expression on lymphocytes can be induced upon activation and TGF- β stimulation. In association with integrin β 7, CD103 is expressed as α E/ β 7 heterodimer. Mature CD103 protein can be cleaved into 2 chains, a 150 kD (C-terminal) chain and a 25 kD (N-terminal) chain, which remain linked by disulfide bonds. CD103 binds to E-cadherin and mediates

homing of lymphocytes to the intestinal epithelium.

Form: Liquid

Conjugation: Unconjugated

Size: 25µg, 100µg

Host Species: Rat

Isotype: Rat IgG2a, κ

Isotype Control: Purified Rat IgG2a, κ Isotype Control[2A3] [Product AGEL0330]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Shipping: Biological ice pack at 4°C



Stability & Storage: Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to

light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial

contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is \leq 2.0 μ g per 106 cells in 100 μ L volume or 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal

performance for each application.