

Product Datasheet Biotin Anti-Mouse CD103 Antibody [M290] Catalogue Code: AGEL0331

Antibody Data

| Product SKU: | AGEL0331 | Clone: | M290 |
|---------------|----------|--------|------|
| Applications: | FCM | | |
| Reactivity: | Mouse | | |

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

| Alternate Names: | Integrin alpha-E;Itgae;Integrin alpha M290;CD103; |
|------------------|---|
| Uniprot ID: | Q60677 |
| Background: | CD103 is a type I transmembrane glycoprotein known as α E integrin or Integrin α IEL chain. It belongs to the integrin family and is primarily found on intestinal intraepithelial lymphocytes (IEL). CD103 is also expressed on a subpopulation of lamina propria T cells, epithelial dendritic cells, lamina propria-derived dendritic cells, and a small subset of peripheral lymphocytes. T regulatory cells express high level of CD103. The CD103 expression on lymphocytes can be induced upon activation and TGF- β stimulation. In association with integrin β 7, CD103 is expressed as α E/ β 7 heterodimer. Mature CD103 protein can be cleaved into 2 chains, a 150 kD (C-terminal) chain and a 25 kD (N-terminal) chain, which remain linked by disulfide bonds. CD103 binds to E-cadherin and mediates homing of lymphocytes to the intestinal epithelium. |
| Form: | Liquid |
| Conjugation: | Biotin |
| Size: | 25µg, 100µg |
| Host Species: | Rat |
| Isotype: | Rat IgG2a, κ |

| Isotype Control: | Biotin Rat IgG2a, κ Isotype Control[2A3] [Product AGEL0331] |
|------------------|---|
| Storage Buffer: | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant. |
| Shipping: | Biological ice pack at 4°C |



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \ \mu$ g per 106 cells in 100 μ L volume or 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.