

Product Datasheet

Biotin Anti-Mouse CD8a Antibody [53-6.7]

Catalogue Code: AGEL0379

Antibody Data

Product SKU: AGEL0379 Clone: 53-6.7

Applications: FCM

Reactivity: Mouse

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: T-cell surface glycoprotein CD8 alpha chain;CD8A;T-lymphocyte differentiation antigen

T8/Leu-2;MAL;

Uniprot ID: P01731

Background: CD8, also known as Lyt-2, Ly-2, or T8, consists of disulfide-linked α and β chains that form

the $\alpha(CD8a)/\beta(CD8b)$ heterodimer and α/α homodimer. CD8a is a 34 kD protein that belongs to the immunoglobulin family. The CD8 α/β heterodimer is expressed on the surface of most thymocytes and a subset of mature TCR α/β T cells. CD8 expression on mature T cells is non-overlapping with CD4. The CD8 α/α homodimer is expressed on a subset of γ/δ TCR-bearing T cells, NK cells, intestinal intraepithelial lymphocytes, and lymphoid dendritic cells. CD8 is an antigen co-receptor on T cells that interacts with MHC class I on antigen-presenting cells or epithelial cells. CD8 promotes T cell activation

through its association with the TCR complex and protein tyrosine kinase lck.

Form: Liquid

Conjugation: Biotin

Size: 25µg, 100µg

Host Species: Rat

Isotype: Rat IgG2a, κ

Isotype Control: Biotin Rat IgG2a, κ Isotype Control[2A3] [Product AGEL0379]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



Stability & Storage: Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to

light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial

contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is \leq 1.0 μ g per 106 cells in 100 μ L volume or 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.