



Product Datasheet

Low Endotoxin Purified Anti-Mouse CD80 Antibody [16-10A1]

Catalogue Code: AGEL0523

Antibody Data

Product SKU:	AGEL0523	Clone:	16-10A1
Applications:	FCM;Block		
Reactivity:	Mouse		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	T-lymphocyte activation antigen CD80;Cd80;Activation B7-1 antigen;B7;
Uniprot ID:	Q00609
Background:	CD80 is a 60 kD highly glycosylated protein. It is a member of the Ig superfamily and is also known as B7-1, B7, and Ly-53. CD80 is constitutively expressed on dendritic cells and monocytes/macrophages, and inducibly expressed on activated B and T cells. The ligation of CD28 on T cells with CD80 and CD86 (B7-2) on antigen presenting cells (such as dendritic cells, macrophages, and B cells) elicits co-stimulation of T cells resulting in enhanced cell activation, proliferation, and cytokine production. CD80 appears to be expressed later in the immune response than CD86. CD80 can also bind to CD152, also known as CTLA-4, to deliver an inhibitory signal to T cells.
Form:	Liquid
Conjugation:	None (AF/LE)
Size:	50µg, 500µg, 1mg
Host Species:	Armenian Hamster
Isotype:	Armenian Hamster IgG
Isotype Control:	AF/LE Purified Armenian Hamster IgG Isotype Control[PIP] [Product AGEL0523]
Storage Buffer:	0.2 µm filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers or stabilizers. Endotoxin level is < 2 EU/mg as Determined by LAL gel clotting assay.
Shipping:	Biological ice pack at 4°C

Stability & Storage: Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

Recommended Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \mu\text{g}$ per 10^6 cells in 100 μL volume or 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
