

Product Datasheet **PE/Cyanine7 Anti-Mouse CD80 Antibody** [16-10A1] Catalogue Code: AGEL0536

Antibody Data

Product SKU:	AGEL0536	Clone:	16-10A1
Applications:	FCM		
Reactivity:	Mouse		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: Uniprot ID:	T-lymphocyte activation antigen CD80;Cd80;Activation B7-1 antigen;B7; Q00609		
Background:	CD80 is a 60 kD highly glycosylated protein. It is a member of the Ig superfamily and is also known as B7-1, B7, and Ly-53. CD80 is constitutively expressed on dendritic cells and monocytes/macrophages, and inducibly expressed on activated B and T cells. The ligation of CD28 on T cells with CD80 and CD86 (B7-2) on antigen presenting cells (such as dendritic cells, macrophages, and B cells) elicits co-stimulation of T cells resulting in enhanced cell activation, proliferation, and cytokine production. CD80 appears to be expressed later in the immune response than CD86. CD80 can also bind to CD152, also known as CTLA-4, to deliver an inhibitory signal to T cells.		
Form:	Liquid	PE/Cyanine7 Excitation and Emission Spectra	
Conjugation:	PE/Cyanine 7	100 -	
Size:	25µg, 100µg		
Host Species:	Armenian Hamster	40	
Isotype:	Armenian Hamster IgG	20 0 350 400 450 500 550 550 550 550 600 650 750 750 800 850 Wavelength (nm) Ex:495;565;755 nm; Em:775 nm	

Isotype Control: PE/Cyanine7 Armenian Hamster IgG Isotype Control[PIP] [Product AGEL0536]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- RecommendedEach lot of this antibody is quality control tested by flow cytometric analysis. Please check
your vial before the experiment. Since applications vary, the appropriate dilutions must be
determined for individual use. We suggest each investigator should titrate the reagent to
obtain optimal results [The recommended concentration is 0.1-1 μg/106 cells in 100 μL
volume].