



## Product Datasheet

### Low Endotoxin Purified Anti-Mouse CD62L Antibody [Mel14]

Catalogue Code: AGEL0626

#### Antibody Data

<b>Product SKU:</b>	<b>AGEL0626</b>	<b>Clone:</b>	<b>Mel14</b>
<b>Applications:</b>	<b>FCM;Neut</b>		
<b>Reactivity:</b>	<b>Mouse</b>		

#### Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

#### Product Information:

<b>Alternate Names:</b>	L-selectin;Sell;CD62 antigen-like family member L;LAM-1;LECAM1;Lymph node homing receptor;Ly-22; CD62L;Lnhr;Ly22;
<b>Uniprot ID:</b>	P18337
<b>Background:</b>	CD62L is a 74-95 kD glycoprotein also known as L-selectin, LECAM-1, Ly-22, LAM-1, and MEL-14. It is a member of the selectin family and is expressed on the majority of B and naïve T cells, a subset of memory T cells, monocytes, granulocytes, most thymocytes, and a subset of NK cells. CD62L is important in lymphocyte homing to high endothelial venules (HEV) in peripheral lymph nodes and leukocyte 'rolling' on activated endothelium. CD62L also contributes to neutrophil emigration at inflammatory sites. CD62L is rapidly shed from lymphocytes and neutrophils upon cellular activation and the expression levels of CD62L (in conjunction with other markers) have been used to distinguish naïve, effector, and memory T cells. CD62L has been reported to interact with CD34, GlyCAM-1, and MAdCAM-1.
<b>Form:</b>	Liquid
<b>Conjugation:</b>	None (AF/LE)
<b>Size:</b>	50µg, 500µg, 1mg
<b>Host Species:</b>	Rat
<b>Isotype:</b>	Rat IgG2a, κ
<b>Isotype Control:</b>	AF/LE Purified Rat IgG2a, κ Isotype Control[2A3] [Product AGEL0626]
<b>Storage Buffer:</b>	0.2 µm filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers or stabilizers. Endotoxin level is < 2 EU/mg as Determined by LAL gel clotting assay.
<b>Shipping:</b>	Biological ice pack at 4°C

**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

**Recommended Usage:** Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 0.25 \mu\text{g}$  per  $10^6$  cells in 100  $\mu\text{L}$  volume or 100  $\mu\text{L}$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.