

Product Datasheet **PE/Cyanine5.5 Anti-Human CD86 Antibody [BU63]** Catalogue Code: AGEL0649

Antibody Data

Product SKU:	AGEL0649	Clone:	BU63
Applications:	FCM		
Reactivity:	Human		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: Uniprot ID: Background:	T-lymphocyte activation antigen CD86;Cd86;Activation B7-2 antigen;Early T-cell costimulatory molecule 1;ETC-1; P42081 CD86 is an 80 kD immunoglobulin superfamily member also known as B7-2, B70, and Ly-58. CD86 is expressed on activated B and T cells, monocytes/macrophages, dendritic cells, and astrocytes. CD86, along with CD80, is the ligand of CD28 and CD152 (CTLA-4). CD86 is expressed earlier in the immune response than CD80. CD86 has also been shown to be involved in immunoglobulin class-switching and triggering of NK cell-mediated cytotoxicity. CD86 binds to CD28 to transduce costimulatory signals for T cell activation, proliferation, and cytokine production. CD86 can bind to CD152 as well, also known as CTLA-4, to deliver an inhibitory signal to T cells.		
Form:	Liquid	PE/Cyanine5.5 Excitation and Emission Spectra	
Conjugation:	PE/Cyanine 5.5		
Size:	20 Tests, 100 Tests, 200 Tests	80 - 87 - 8 60 -	
Host Species:	Mouse	S 60	
Isotype:	Mouse IgG1, κ	20 0 350 400 450 500 500 500 500 500 600 650 700 750 800 850 850 850 850 850 850 8	
Isotype Control:	PE/Cyanine5 5 Mouse IgG1 x Isotype Control[MOPC-21] [Product AGE] 0640]		

Isotype Control: PE/Cyanine5.5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL0649]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.