

Product Datasheet

PerCP Anti-Human/Mouse/Rat CD47 Antibody [MIAP410] Catalogue Code: AGEL0672

Antibody Data


Product SKU:	AGEL0672	Clone:	MIAP410
Applications:	FCM		
Reactivity:	Human;Mouse;Rat		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: Uniprot ID: Background:	Leukocyte surface antigen CD47;Cd47;Integrin-associated protein;IAP; Q08722 Q61735 P97829 CD47, also known as Integrin-Associated Protein (IAP), is a membrane protein of about 50 kD with an IgV-like extracelluluar domain, a five membrane-spanning segment and a short terminal cytoplasmic region. It is widely expressed on many cell types and often associated with beta 3 integrins. It has been reported that CD47 functions as a self marker. Red cells lacking CD47 were rapidly cleared from the bloodstream by splenic macrophages. By binding to SIRPα, CD47 controls hemostatic innate immune functions, such as phagocytosis and cell trafficking.		
Form:	Liquid	PerCP Excitation and Emission Spectra	
Conjugation:	PerCP	100 -	
Size:	20 Tests, 100 Tests, 200 Tests	80 3 5 60 	
Host Species:	Mouse	€ 60 100 11 10 10 10 10 10 10 10 10 10 10 1	
Isotype:	Mouse IgG1, к	20 0 350 400 450 500 550 600 650 700 750 800 Wavelength (nm)	



Isotype Control: PerCP Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL0672]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.