

Product Datasheet

PE/Cyanine7 Anti-Human/Mouse/Rat CD47 Antibody [MIAP410] Catalogue Code: AGEL0684

Antibody Data

Product SKU:	AGEL0684	Clone:	MIAP410
Applications:	FCM		
Reactivity:	Human;Mouse;Rat		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	Leukocyte surface antigen CD47;Cd47;Integrin-associated protein;IAP;		
Uniprot ID:	Q08722 Q61735 P97829		
Background:	CD47, also known as Integrin-Associated Protein (IAP), is a membrane protein of about 50 kD with an IgV-like extracelluluar domain, a five membrane-spanning segment and a short terminal cytoplasmic region. It is widely expressed on many cell types and often associated with beta 3 integrins. It has been reported that CD47 functions as a self marker. Red cells lacking CD47 were rapidly cleared from the bloodstream by splenic macrophages. By binding to SIRP α , CD47 controls hemostatic innate immune functions, such as phagocytosis and cell trafficking.		
Form:	Liquid	PE/Cyanine7 Excitation and Emission Spectra	
Conjugation:	PE/Cyanine 7	100 -	
Size:	25µg, 100µg		
Host Species:	Mouse	40	
Isotype:	Mouse IgG1, к	20 0 350 400 450 500 550 600 650 700 750 800 850 Wavelength (nm)	

Ex:495;565;755 nm; Em:775 nm

Isotype Control: PE/Cyanine7 Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL0684]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- RecommendedEach lot of this antibody is quality control tested by flow cytometric analysis. Please check
your vial before the experiment. Since applications vary, the appropriate dilutions must be
determined for individual use. We suggest each investigator should titrate the reagent to
obtain optimal results [The recommended concentration is 0.1-1 μg/106 cells in 100 μL
volume].