

Product Datasheet **PE/Cyanine5 Anti-Mouse CD127/IL-7RA Antibody [A7R34]** Catalogue Code: AGEL0740

Antibody Data

Product SKU:	AGEL0740	Clone:	A7R34
Applications:	FCM		
Reactivity:	Mouse		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: Uniprot ID:	Interleukin-7 receptor subunit alpha;II7r;IL-7 receptor subunit alpha;IL-7R subunit alpha;IL- 7R-alpha;IL-7RA;CD127; P16872		
Background:	CD127 is a 60-90 kD type I transmembrane glycoprotein also known as IL-7 receptor α chain or IL-7Rα. It forms a heterodimer with the common γ chain (γc or CD132) which is shared with the receptors for IL-2, IL-4, IL-9, IL-13, IL-15, and IL-21. CD127 is expressed on immature B cells through early pre-B stage, thymocytes (except CD4/CD8 double positive thymocytes), peripheral T cells, and bone marrow stromal cells. CD127 has been reported to be an useful marker for identifying memory and effector T cells. The ligation of IL-7 with its receptor is important for stimulation of mature and immature T cells as well as immature B cells proliferation and development.		
Form:	Liquid	PE/Cyanine5 Excitation and Emission Spectra	
Conjugation:	PE/Cyanine 5	100 -	
Size:	50 Tests, 100 Tests, 200 Tests	80 - 37 - 17 - 60 -	
Host Species:	Rat	40	
Isotype:	Rat IgG2a, κ	20 0 350 400 450 500 550 500 550 600 650 700 750 800 Wavelength (nm)	
		Ex:495;565;655 nm; Em:670 nm	
Isotype Control:	PE/Cyanine5 Rat IgG2a, K Isotyne Control[2A3] [Product AGEI 0740]		

Isotype Control: PE/Cyanine5 Rat IgG2a, κ Isotype Control[2A3] [Product AGEL0740]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.