

Product Datasheet **PE Anti-Mouse CD127/IL-7RA Antibody [A7R34]** Catalogue Code: AGEL0746

Antibody Data

Product SKU:	AGEL0746	Clone:	A7R34
Applications:	FCM		
Reactivity:	Mouse		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	Interleukin-7 receptor subunit alpha;II7r;IL-7 receptor subunit alpha;IL-7R subunit alpha;IL- 7R-alpha;IL-7RA;CD127;		
Uniprot ID: Background:	P16872 CD127 is a 60-90 kD type I transmembrane glycoprotein also known as IL-7 receptor α chain or IL-7Rα. It forms a heterodimer with the common γ chain (γc or CD132) which is shared with the receptors for IL-2, IL-4, IL-9, IL-13, IL-15, and IL-21. CD127 is expressed on immature B cells through early pre-B stage, thymocytes (except CD4/CD8 double positive thymocytes), peripheral T cells, and bone marrow stromal cells. CD127 has been reported to be an useful marker for identifying memory and effector T cells. The ligation of IL-7 with its receptor is important for stimulation of mature and immature T cells as well as immature B cells proliferation and development.		
Form:	Liquid	PE Excitation and Emission Spectra	
Conjugation:	PE	100	
Size:	25µg, 100µg	80 - (\$) 19 60 -	
Host Species:	Rat	(%) 500	
Isotype:	Rat IgG2a, к	20 0 350 400 450 500 550 600 650 700 Wavelength (nm)	
		Ex:495;565 nm; Em:575 nm	
Isotupe Control:	PE Pat Jac2a, K Jaatura Control[2A2] [Product ACEL 0746]		

Isotype Control: PE Rat IgG2a, κ Isotype Control[2A3] [Product AGEL0746]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- RecommendedEach lot of this antibody is quality control tested by flow cytometric analysis. Please check
your vial before the experiment. Since applications vary, the appropriate dilutions must be
determined for individual use. We suggest each investigator should titrate the reagent to
obtain optimal results [The recommended concentration is 0.1-1 μg/106 cells in 100 μL
volume].