

Product Datasheet **FITC Anti-Mouse CD122 Antibody [5H4]** Catalogue Code: AGEL0791

## Antibody Data

Applications: FCM	Product SKU:	AGEL0791	Clone:	5H4	
	Applications:	FCM			
Reactivity: Mouse	Reactivity:	Mouse			

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

## Product Information:

Alternate Names: Uniprot ID:	Interleukin-2 receptor subunit beta;II2rb;IL-2 receptor subunit beta; IL-2R subunit beta;IL- 2RB;High affinity IL-2 receptor subunit beta;p70-75;CD122; P16297		
Background:	CD122 is a 70-75 kD IL-2 receptor $\beta$ chain also known as IL-2R $\beta$ , which is also shared by the IL-15 receptor. It is constitutively expressed by NK cells and at lower levels by T cells, B cells, monocytes, and macrophages. The IL-2R $\beta$ chain can combine with either the common $\gamma$ subunit ( $\gamma$ c, CD132) alone or with the $\gamma$ c subunit and the IL-2R $\alpha$ subunit (CD25) to generate intermediate or high affinity IL-2 receptor complexes, respectively. CD122 expression levels can be upregulated by activation. The 5H4 antibody does not block IL-2 binding to the IL-2 receptor. CD122 is expressed on murine, but not human, CD8+ Tregs involved in the maintenance of T cell homeostasis.		
Form:	Liquid	FITC Excitation and Emission Spectra	
Conjugation:	FITC	100 -	
		80 -	
Size:	50 Tests, 100 Tests, 200 Tests	80 - 8 60 -	
Size: Host Species:	50 Tests, 100 Tests, 200 Tests Rat	2	



**Isotype Control:** FITC Rat IgG2a, κ Isotype Control[2A3] [Product AGEL0791]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.