

Product Datasheet

APC Anti-Mouse CD122 Antibody [5H4]

Catalogue Code: AGEL0801

Antibody Data

Product SKU: AGEL0801 Clone: 5H4

Applications: FCM

Reactivity: Mouse

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: Interleukin-2 receptor subunit beta; IL-2R subuni

2RB;High affinity IL-2 receptor subunit beta;p70-75;CD122;

Uniprot ID: P16297

Background: CD122 is a 70-75 kD IL-2 receptor β chain also known as IL-2R β , which is also shared by

the IL-15 receptor. It is constitutively expressed by NK cells and at lower levels by T cells, B cells, monocytes, and macrophages. The IL-2R β chain can combine with either the common γ subunit (γ c, CD132) alone or with the γ c subunit and the IL-2R α subunit (CD25) to generate intermediate or high affinity IL-2 receptor complexes, respectively. CD122 expression levels can be upregulated by activation. The 5H4 antibody does not block IL-2 binding to the IL-2 receptor. CD122 is expressed on murine, but not human, CD8+ Tregs

involved in the maintenance of T cell homeostasis.


Form: Liquid

Conjugation: APC

Size: 25µg, 100µg

Host Species: Rat

Isotype: Rat IgG2a, κ



Ex:650 nm; Em:660 nm

Isotype Control: APC Rat IgG2a, κ Isotype Control[2A3] [Product AGEL0801]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



Stability & Storage: Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to

light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial

contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/106 cells in 100 μ L

volume].