

GenieFluor 647 Anti-Mouse CD122 Antibody [5H4]
 Catalogue Code: AGEL0807

Antibody Data

Product SKU:	AGEL0807	Clone:	5H4
Applications:	FCM		
Reactivity:	Mouse		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.


Product Information:

Alternate Names: Interleukin-2 receptor subunit beta;Il2rb;IL-2 receptor subunit beta; IL-2R subunit beta;IL-2RB;High affinity IL-2 receptor subunit beta;p70-75;CD122;
Uniprot ID: P16297

Background: CD122 is a 70-75 kD IL-2 receptor β chain also known as IL-2R β , which is also shared by the IL-15 receptor. It is constitutively expressed by NK cells and at lower levels by T cells, B cells, monocytes, and macrophages. The IL-2R β chain can combine with either the common γ subunit (γ c, CD132) alone or with the γ c subunit and the IL-2R α subunit (CD25) to generate intermediate or high affinity IL-2 receptor complexes, respectively. CD122 expression levels can be upregulated by activation. The 5H4 antibody does not block IL-2 binding to the IL-2 receptor. CD122 is expressed on murine, but not human, CD8+ Tregs involved in the maintenance of T cell homeostasis.

Form: Liquid
Conjugation: Genie Fluor647
Size: 25 μ g, 100 μ g
Host Species: Rat
Isotype: Rat IgG2a, κ

647 Excitation and Emission Spectra



Isotype Control: Genie Fluor 647 Rat IgG2a, κ Isotype Control[2A3] [Product AGEL0807]
Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Shipping: Biological ice pack at 4°C

Stability & Storage: Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

Recommended Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 µg/10⁶ cells in 100 µL volume].