

Product Datasheet **PerCP/Cyanine5.5 Anti-Human CD5 Antibody [UCHT2]** Catalogue Code: AGEL0893

Antibody Data

Product SKU:	AGEL0893	Clone:	UCHT2	
Applications:	FCM			
Reactivity:	Human			

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	T-cell surface glycoprotein CD5;CD5;Lymphocyte antigen T1/Leu-1;LEU1;		
Uniprot ID:	P06127		
Background:	CD5 is a 67 kD single chain type I glycoprotein also known as Leu-1, Ly-1 and T1. It is a member of the scavenger receptor superfamily found on T cells, thymocytes, B cell subsets, chronic B lymphocytic leukemia (B-Cells), and peripheral blood dendritic cells. CD5 modulates T and B cell receptor signaling, thymocyte maturation, and T-B cell interactions upon binding to ligands such as CD72.		
Form:	Liquid	PerCP/Cyanine5.5 Excitation and Emission Spectra	
Conjugation:	PerCP/Cyanine 5.5	100	
Size:	20 Tests, 100 Tests, 200 Tests		
Host Species:	Mouse	€) pp 60 10 10 10	
Isotype:	Mouse IgG1, к	20 0 350 400 450 500 550 600 650 700 750 800 850 Wavelength (nm)	
		Ex:440;480;675 nm; Em:675 nm	
Isotype Control:	PerCP/Cyanine5 5 Mouse IgG1 r Isotype Control[MOPC-21] [Product AGEL0893]		

Isotype Control: PerCP/Cyanine5.5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL0893]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.