

Product Datasheet GenieFluor 647 Anti-Human CD45RA Antibody [HI100]

Catalogue Code: AGEL0961

## Antibody Data

Product SKU:	AGEL0961	Clone:	HI100	
Applications:	FCM			
Reactivity:	Human			

## Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

## **Product Information:**

Alternate Names:	Receptor-type tyrosine-protein phosphatase C;Ptprc;L-CA;Ly-5;T200;CD45;		
Uniprot ID:	P08575		
Background:	CD45RA is a 205-220 kD single chain type I glycoprotein. It is an exon 4 splice variant of the tyrosine phosphatase CD45. The CD45RA isoform is expressed on resting/naïve T cells, medullary thymocytes, B cells and monocytes. CD45RA enhances both T cell receptor and B cell receptor signaling. CD45 non-covalently associates with lymphocyte phosphatase-associated phosphoprotein (LPAP) on T and B lymphocytes. CD45 has been reported to be associated with several other cell surface antigens including CD1, CD2, CD3, and CD4. CD45 has also been reported to bind galectin-1. CD45 isoform expression can change in response to cytokines.		
Form:	Liquid	647 Excitation and Emission Spectra	
Conjugation:	Genie Fluor647	100 -	
Size:	20 Tests, 100 Tests, 200 Tests	80 - S	
Host Species:	Mouse	(%) po	
Isotype:	Mouse IgG2b, к	E 40 20 0 350 400 450 500 550 600 650 700 750 800 850 Wavelength (nm)	
Isotype Control:	Genie Fluor 647 Mouse IgG2b, κ Isotype Control[MPC-11] [Product AGEL0961]		

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.