

Product Datasheet **FITC Anti-Human CD11a Antibody [TS1/22.1.1.13]** Catalogue Code: AGEL0962

Antibody Data

Product SKU:	AGEL0962	Clone:	TS1/22.1.1.13
Applications:	FCM		
Reactivity:	Human		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: Uniprot ID:	Integrin alpha-L;Itgal;CD11 antigen-like family member A;LFA-1A;Lymphocyte antigen 15;Ly-15;CD11a; P20701		
Background:	CD11a is a 170-180 kD type I transmembrane glycoprotein also known as LFA-1 α chain and integrin α L subunit. CD11a non-covalently associates with integrin β 2 (CD18) to form LFA-1. It is expressed on all leukocytes, including B and T lymphocytes, monocytes, macrophages, neutrophils, basophils and eosinophils. It is absent on non-hematopoietic tissues and platelets. CD11a plays a central role in leukocyte cell-cell interactions and is important in lymphocyte costimulation. CD11a/CD18 binds to ICAM-1 (CD54), ICAM-2 (CD102), and ICAM-3 (CD50).		
Form:	Liquid	FITC Excitation and Emission Spectra	
Conjugation:	FITC	80	
Size:	20 Tests, 100 Tests, 200 Tests		
Host Species:	Mouse	60	
Isotype:	Mouse IgG1, κ	20 0 350 400 450 500 550 600 650 700 Wavelength (nm)	



Isotype Control: FITC Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL0962]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.