

Product Datasheet GenieFluor 647 Anti-Human CD38 Antibody [HIT2]

Catalogue Code: AGEL0993

## Antibody Data

Product SKU:	AGEL0993	Clone:	HIT2
Applications:	FCM		
Reactivity:	Human		

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

## **Product Information:**

Alternate Names: Uniprot ID:	ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1;CD38;2'-phospho-ADP-ribosyl cyclase;2'-phospho-cyclic-ADP-ribose transferase;ADP-ribosyl cyclase 1;ADPRC 1;Cyclic ADP-ribose hydrolase 1;cADPr hydrolase 1;T10;CD38; P28907		
Background:	CD38 is a 45 kD type II transmembrane glycoprotein also known as T10. It is an ADP- ribosyl hydrolase expressed at variable levels on hematopoietic cells and in some non- hematopoietic tissues (such as brain, muscles, and kidney). In humans, it is expressed at high levels on plasma cells and activated T and B cells. By functioning as both a cyclase and a hydrolase, CD38 mediates lymphocyte activation, adhesion, and the metabolism of cADPR and NAADP. CD31 is the ligand of CD38.		
Form:	Liquid	647 Excitation and Emission Spectra	
Conjugation:	Genie Fluor647	100 -	
Size:	20 Tests, 100 Tests, 200 Tests	80 -	
Host Species:	Mouse	(%) po po po po po po po po	
Isotype:	Mouse IgG1, κ	E 40 20 0 350 400 450 500 550 600 650 700 750 800 850 Wavelength (nm)	

**Isotype Control:** Genie Fluor 647 Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL0993]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.