

Product Datasheet GenieFluor 488 Anti-Human CD35 Antibody [E11] Catalogue Code: AGEL1015

Antibody Data

Product SKU:	AGEL1015	Clone:	E11
Applications:	FCM		
Reactivity:	Human		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	Complement receptor type 1;CR1;C3b/C4b receptor;CD35;CR1;C3BR;		
Uniprot ID:	P17927		
Background:	CD35 is a type I single chain of glycoprotein, also known as C3b/C4b receptor, Complement Receptor type 1 or CR1. Four molecular weight allotypes (160kD, 190kD, 220kD, and 250kD) have been described. CD35 is expressed on granulocytes, monocytes, B cells, erythrocytes, and follicular dendritic cells, as well as subsets of NK and T cells. CD35 binds complement C3b, C4b, or iC3, and iC4, and plays important roles in both innate and adoptive immune response via mediating phagocytosis by granulocytes and monocytes. CD35 has also been reported to inhibit T-cell proliferation.		
Form:	Liquid	488 Excitation and Emission Spectra	
Conjugation:	Genie Fluor488	100 -	
Size:	20 Tests, 100 Tests, 200 Tests		
Host Species:	Mouse	(%) p ₉ jieu 40 .	
Isotype:	Mouse IgG1, κ	\dot{z} \dot{u}_{0} \dot{z}_{0} $$	

Isotype Control: Genie Fluor 488 Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL1015]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.