

Product Datasheet **PE Anti-Human CD58 Antibody [TS2/9.1]** Catalogue Code: AGEL1038

## Antibody Data


Product SKU:	AGEL1038	Clone:	TS2/9.1
Applications:	FCM		
Reactivity:	Human		

## Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

## **Product Information:**

Alternate Names:	Lymphocyte function-associated antigen 3;CD58;Ag3;Surface glycoprotein LFA-3;LFA3;		
Uniprot ID:	P19256		
Background:	CD58, also known as lymphocyte function-associated antigen 3 (LFA-3) is a 45-70 kD cell surface protein that is a member of the immunoglobulin superfamily. Alternative splicing of CD58 gives rise to transmembrane and glycosylphosphatidylinositol (GPI)-anchored forms on cell surface. CD58 is expressed on both hematopoietic and non-hematopoietic cells including B cells, T cells, monocytes, erythrocytes, endothelial cells, epithelial cells, and fibroblasts. High levels are observed on memory T cells and dendritic cells. CD58 expressed on antigen presenting cells and target cells enhances T cell recognition via the binding of it's cognate ligand, CD2, on the T cell surface.		
Form:	Liquid	PE Excitation and Emission Spectra	
Conjugation:	PE	100	
Size:	20 Tests, 100 Tests, 200 Tests	- 08 	
Host Species:	Mouse	60 50	
Isotype:	Mouse IgG1, к	20 0 350 400 450 550 550 550 550 550 550 550 5	



**Isotype Control:** PE Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL1038]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.