

Product Datasheet **PE/Cyanine5 Anti-Human CD58 Antibody [TS2/9.1]** Catalogue Code: AGEL1040

Antibody Data

Applications:FCMReactivity:Human	Product SKU:	AGEL1040	Clone:	TS2/9.1	
Reactivity: Human	Applications:	FCM			
	Reactivity:	Human			

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	Lymphocyte function-associated antigen 3;CD58;Ag3;Surface glycoprotein LFA-3;LFA3;		
Uniprot ID:	P19256		
Background:	CD58, also known as lymphocyte function-associated antigen 3 (LFA-3) is a 45-70 kD cell surface protein that is a member of the immunoglobulin superfamily. Alternative splicing of CD58 gives rise to transmembrane and glycosylphosphatidylinositol (GPI)-anchored forms on cell surface. CD58 is expressed on both hematopoietic and non-hematopoietic cells including B cells, T cells, monocytes, erythrocytes, endothelial cells, epithelial cells, and fibroblasts. High levels are observed on memory T cells and dendritic cells. CD58 expressed on antigen presenting cells and target cells enhances T cell recognition via the binding of it's cognate ligand, CD2, on the T cell surface.		
Form:	Liquid	PE/Cyanine5 Excitation and Emission Spectra	
Conjugation:	PE/Cyanine 5	100 -	
Size:	20 Tests, 100 Tests, 200 Tests	8 00 00 00 00 00 00 00 00 00 00 00 00 00	
Host Species:	Mouse	Те 40 - / / / / / / / / / / / / / / / / / /	
Isotype:	Mouse IgG1, к	20 0 350 400 450 500 550 600 650 700 750 800	
		Wavelength (nm)	
		Ex:495;565;655 nm; Em:670 nm	
Isotype Control:	-		
Storage Buffer:	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.		

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.