

Product Datasheet GenieFluor 488 Anti-Human CD235 Antibody [HIR2]

Catalogue Code: AGEL1097

## Antibody Data

Product SKU:	AGEL1097	Clone:	HIR2
Applications:	FCM		
Reactivity:	Human		

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

## **Product Information:**

Alternate Names: Uniprot ID:	Glycophorin-A/B;GYPA/B;MN sialoglycoprotein;SS-active sialoglycoprotein;PAS- 2/3;Sialoglycoprotein alpha/delta;CD235a/b; P02724 P06028		
Background:	The HIR2 antibody reacts with a common epitope of glycophorin A (CD235a) and glycophorin B (CD235b). Glycophorin A is the major sialoglycoprotein expressed on red blood cell membrane, and erythroid precursors. Glycophorin A shares strong homology with glycophorin B. The HIR2 antibody recognizes human RBCs and erythroid precursors and is useful in erythroid cell development studies. Mature, non-nucleated red blood cells are characteristically glycophorin A positive, but CD45 and CD71 negative.		
Form:	Liquid	488 Excitation and Emission Spectra	
Conjugation:	Genie Fluor488	100	
Size:	20 Tests, 100 Tests, 200 Tests	80 - S	
Host Species:	Mouse	(%) 60	
Isotype:	Mouse IgG2b, κ	20 0 350 400 450 500 550 600 650 700 Wavelength (nm)	

Isotype Control: Genie Fluor 488 Mouse IgG2b, κ Isotype Control[MPC-11] [Product AGEL1097]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.