

Product Datasheet **APC Anti-Human CD41 Antibody [HIP8]** Catalogue Code: AGEL1132

Antibody Data

| Product SKU: | AGEL1132 | Clone: | HIP8 |
|---------------|----------|--------|------|
| Applications: | FCM | | |
| Reactivity: | Human | | |

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

| Alternate Names: Uniprot ID: | Integrin alpha-Iib;ITGA2B;GPal Iib;CD41;GP2B; ITGAB; P08514 | oha lib;GPIIb;Platelet membrane glycoprotein | |
|---------------------------------|--|---|--|
| Background: | CD41 is a 125/25 kD α subunit of the gpIIb/IIIa (CD41/CD61) complex. CD41 is a heterodimer composed of a heavy chain (gpIIb α) and light chain (gpIIb β) linked by a single disulfide bond. It is a member of the integrin family primarily expressed on platelets and megakaryocytes. CD41 has been reported to be involved with platelet aggregation and platelet attachment to the ECM. CD41/CD61 complex acts as the receptor for fibrinogen, fibronectin, Von Willebrand factor and thrombin. | | |
| Form: | Liquid | APC Excitation and Emission Spectra | |
| Conjugation: | APC | 80 - | |
| Size: | 20 Tests, 100 Tests, 200 Tests | | |
| Host Species: | Mouse | 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 | |
| Isotype: | Mouse IgG1, к | 20 0 350 400 450 500 550 600 650 700 750 800 850 Wavelength (nm) | |
| | | Ex:650 nm; Em:660 nm | |

Isotype Control: APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL1132]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.