

Product Datasheet

Low Endotoxin Purified Anti-Mouse CD90.2 Antibody [30H12] Catalogue Code: AGEL1205

## Antibody Data

Product SKU:	AGEL1205	Clone:	30H12
Applications:	FCM;Depletion		
Reactivity:	Mouse		

## Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

## Product Information:

Alternate Names: Uniprot ID: Background:	Thy-1.2 membrane glycoprotein;Thy1.2;Thy-1.2 antigen;CD90.2;Thy-1.2; - CD90.2 is a 25-35 kD immunoglobulin superfamily member also known as Thy1.2. It is expressed on hematopoietic stem cells and neurons, all thymocytes, and peripheral T cells in Thy1.2 bearing mouse strains (Balb/c, CBA/J, C3H/He, C57BL/-, DBA, NZB/-). CD90.2 is a glycosylphosphatidylinositol (GPI)-anchored membrane glycoprotein involved in signal transduction. CD90.2 is involved in costimulation of lymphocyte proliferation and induction of hematopoietic stem cells differentiation. CD90.2 has been shown to interact with CD45. The 30H12 antibody has been reported to induce Ca2+ flux in thymocytes and, in combination with antibody against the CD3/TCR complex, promote thymocyte apoptosis and inhibit CD3-mediated proliferative responses of mature T lymphocytes.		
Form:	Liquid		
Conjugation:	None (AF/LE)		
Size:	50µg, 500µg, 1mg		
Host Species:	Rat		
Isotype:	Rat IgG2b, κ		
Isotype Control:	AF/LE Purified Rat IgG2b, κ Isotype Control[LTF-2] [Product AGEL1205]		
Storage Buffer:	0.2 μm filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers or stabilizers. Endotoxin level is < 2 EU/mg as Determined by LAL gel clotting assay.		

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 2.0 \ \mu$ g per 106 cells in 100  $\mu$ L volume or 100  $\mu$ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.