

Product Datasheet **PerCP/Cyanine5.5 Anti-Human/Mouse CD44 Antibody [IM7]** Catalogue Code: AGEL1287

## Antibody Data

Product SKU:	AGEL1287	Clone:	IM7
Applications:	FCM		
Reactivity:	Human;Mouse		

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

## Product Information:

Alternate Names: Uniprot ID:	CD44 antigen;CD44;CDw44;Epican;Phagocytic glycoprotein 1;PGP-1;Phagocytic glycoprotein I;PGP-I;CD44;LHR; MDU2; MDU3; MIC4; P15379 P16070		
Background:	CD44 is a 80-95 kD glycoprotein also known as Hermes, Pgp1, H-CAM, or HUTCH. It is expressed on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. As B and T cells become activated or progress to the memory stage, CD44 expression increases from low or mid levels to high levels. Thus, CD44 has been reported to be a valuable marker for memory cell subsets. High CD44 expression on Treg cells has been associated with potent suppressive function via high production of IL-10. CD44 is an adhesion molecule involved in leukocyte attachment to and rolling on endothelial cells, homing to peripheral lymphoid organs and to the sites of inflammation, and leukocyte aggregation.		
Form:	Liquid	PerCP/Cyanine5.5 Excitation and Emission Spectra	
Conjugation:	PerCP/Cyanine 5.5	100	
Size:	25µg, 100µg		
Host Species:	Rat	Vorumality of the second secon	
Isotype:	Rat IgG2b, κ	20 0 350 400 450 500 550 600 650 700 750 800 850 Wavelength (nm) Ex:440;480;675 nm; Em:675 nm	

Isotype Control: PerCP/Cyanine5.5 Rat IgG2b, κ Isotype Control[LTF-2] [Product AGEL1287]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- RecommendedEach lot of this antibody is quality control tested by flow cytometric analysis. Please check<br/>your vial before the experiment. Since applications vary, the appropriate dilutions must be<br/>determined for individual use. We suggest each investigator should titrate the reagent to<br/>obtain optimal results [The recommended concentration is 0.1-1 μg/106 cells in 100 μL<br/>volume].