

**Antibody Data**

|                      |                 |               |                 |
|----------------------|-----------------|---------------|-----------------|
| <b>Product SKU:</b>  | <b>AGEL1321</b> | <b>Clone:</b> | <b>145-2C11</b> |
| <b>Applications:</b> | <b>FCM</b>      |               |                 |
| <b>Reactivity:</b>   | <b>Mouse</b>    |               |                 |


**Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

**Product Information:**

**Alternate Names:** T-cell surface glycoprotein CD3 epsilon chain;CD3E;T-cell surface antigen T3/Leu-4 epsilon chain;CD3e;CD3E;T3E;  
**Uniprot ID:** P22646  
**Background:** CD3ε is a 20 kD transmembrane protein, also known as CD3 or T3. It is a member of the Ig superfamily and primarily expressed on T cells, NK-T cells, and at different levels on thymocytes during T cell differentiation. CD3ε forms a TCR complex by associating with the CD3δ, γ and ζ chains, as well as the TCR α/β or γ/δ chains. CD3 plays a critical role in TCR signal transduction, T cell activation, and antigen recognition by binding the peptide/MHC antigen complex.

**Form:** Liquid  
**Conjugation:** Genie Fluor647  
**Size:** 50 Tests, 100 Tests, 200 Tests  
**Host Species:** Armenian Hamster  
**Isotype:** Armenian Hamster IgG



**Isotype Control:** Genie Fluor 647 Armenian Hamster IgG Isotype Control[PIP] [Product AGEL1321]  
**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.  
**Shipping:** Biological ice pack at 4°C

**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

**Recommended Usage:** Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.