

Product Datasheet APC Anti-Mouse CD45R/B220 Antibody [RA3.3A 1/6.1] Catalogue Code: AGEL1410

Antibody Data

Product SKU:	AGEL1410	Clone:	RA3.3A 1/6.1
Applications:	FCM		
Reactivity:	Mouse		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	B220;		
Uniprot ID:	-		
Background:	CD45R, also known as B220, is an isoform of CD45. It is a member of the protein tyrosine phosphatase (PTP) family with a molecular weight of approximately 180-240 kD. CD45R is expressed on B cells (at all developmental stages from pro-B cells through mature B cells), activated B cells, and subsets of T and NK cells. CD45R (B220) is also expressed on a subset of abnormal T cells involved in the pathogenesis of systemic autoimmunity in MRL-Fas ^{lpr} and MRL-Fas ^{gld} mice. It plays a critical role in TCR and BCR signaling. The primary ligands for CD45 are galectin-1, CD2, CD3, and CD4. CD45R is commonly used as a pan-B cell marker; however, CD19 may be more appropriate for B cell specificity.		
Form:	Liquid	APC Excitation and Emission Spectra	
Conjugation:	APC	100 - 80 -	
Size:	25µg, 100µg		
Host Species:	Rat	Normalized (%)	
Isotype:	Rat IgM, к	20 -	
		350 400 450 500 550 600 650 700 750 800 850 Wavelength (nm)	
		Ex:650 nm; Em:660 nm	
Isotype Control:	APC Rat IgM, κ Isotype Control[RTK2118] [Product AGEL1410]		
Storage Buffer:	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.		

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μg/106 cells in 100 μL volume].