

Product Datasheet GenieFluor 488 Anti-Mouse CD45.2 Antibody [104.2] Catalogue Code: AGEL1533

Antibody Data

Product SKU:	AGEL1533	Clone:	104.2
Applications:	FCM		
Reactivity:	Mouse		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	Ly-5.2; LCA;		
Uniprot ID:	-		
Background:	CD45.2 is an alloantigen of CD45, expressed by Ly5.2 bearing mouse strains (e.g., A, AKR, BALB/c, CBA/Ca, CBA/J, C3H/He, C57BL, C57BR, C57L, C58, DBA/1, DBA/2, NZB, SWR, 129). CD45, a member of the protein tyrosine phosphatase (PTP) family, is a 180-240 kD glycoprotein expressed on all hematopoietic cells except mature erythrocytes and platelets. There are multiple isoforms in the mouse that play key roles in TCR and BCR signal transduction. These isoforms are very specific to the activation and maturation states of the cell as well as specific cell type. The primary ligands for CD45 are galectin-1, CD2, CD3, CD4, TCR, CD22, and Thy-1.		
Form:	Liquid	488 Excitation and Emission Spectra	
Conjugation:	Genie Fluor488	100	
Size:	25µg, 100µg	80 - S	
Host Species:	Mouse	(%) 60	
Isotype:	Mouse IgG2a, к	$\frac{1}{20} \int_{\frac{1}{350}}^{\frac{1}{20}} \frac{1}{400} \int_{\frac{1}{450}}^{\frac{1}{500}} \frac{1}{500} \int_{\frac{1}{500}}^{\frac{1}{500}} \frac{1}{500} \int_{\frac{1}{500}}^{\frac{1}{500}} \frac{1}{650} \int_{\frac{1}{700}}^{\frac{1}{700}} \frac{1}{100} \frac{1}{1$	
Isotype Control:	Genie Fluor 488 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product AGEL1533]		
Storage Buffer:	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.		

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μg/106 cells in 100 μL volume].