

Product Datasheet **Purified Anti-Human CD279/PD-1 Antibody [J116]** Catalogue Code: AGEL1607

Antibody Data

Product SKU:	AGEL1607	Clone:	J116
Applications:	FCM		
Reactivity:	Human		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	PD-1; Programmed Death-1;
Uniprot ID:	Q15116
Background:	Programmed cell death 1 (PD-1), also known as CD279, is a 55 kD member of the immunoglobulin superfamily. CD279 contains the immunoreceptor tyrosine-based inhibitory motif (ITIM) in the cytoplasmic region and plays a key role in peripheral tolerance and autoimmune disease. CD279 is expressed predominantly on activated T cells, B cells, and myeloid cells. PD-L1 (B7-H1) and PD-L2 (B7-DC) are ligands of CD279 (PD-1) and are members of the B7 gene family. Evidence suggests overlapping functions for these two PD-1 ligands and their constitutive expression on some normal tissues and upregulation on activated antigen-presenting cells. Interaction of CD279 ligands results in inhibition of T cell proliferation and cytokine secretion.
Form:	Liquid
Conjugation:	Unconjugated
Size:	25µg, 100µg
Host Species:	Mouse
Isotype:	Mouse IgG1, к

Isotype Control:	Purified Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL1607]
------------------	--------------------------------------------------------------------

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \ \mu$ g per 106 cells in 100 μ L volume or 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.