

Product Datasheet **FITC Anti-Mouse CD279/PD-1 Antibody** [29F.1A12] Catalogue Code: AGEL1627

Antibody Data

Product SKU:	AGEL1627	Clone:	29F.1A12	
Applications:	FCM			
Reactivity:	Mouse			

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Q02242		
CD279, also known as programmed death-1 (PD-1), is a 50-55 kD glycoprotein belonging to the CD28 family of the Ig superfamily. PD-1 is expressed on activated splenic T and B cells and thymocytes. It is induced on activated myeloid cells as well. PD-1 is involved in lymphocyte clonal selection and peripheral tolerance through binding its ligands, B7-H1 (PD-L1) and B7-DC (PD-L2). It has been reported that PD-1 and PD-L1 interactions are critical to positive selection and play a role in shaping the T cell repertoire. PD-L1 negative costimulation is essential for prolonged survival of intratesticular islet allografts.		
Liquid	FITC Excitation and Emission Spectra	
FITC	100	
50 Tests, 100 Tests, 200 Tests		
Rat	60	
Rat IgG2a, к	2 20 350 400 450 500 550 600 600 650 700 Wavelength (nm) Ex:490 nm; Em:530 nm	
	CD279, also known as programmed to the CD28 family of the Ig superfa- cells and thymocytes. It is induced of lymphocyte clonal selection and per (PD-L1) and B7-DC (PD-L2). It has critical to positive selection and play costimulation is essential for prolong Liquid FITC 50 Tests, 100 Tests, 200 Tests Rat	

Isotype Control:	FITC Rat IgG2a, κ Isotype Control[2A3] [Product AGEL1627]	

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.