

Product Datasheet **APC Anti-Mouse CD279/PD-1 Antibody** [29F.1A12] Catalogue Code: AGEL1634

Antibody Data

Product SKU:	AGEL1634	Clone:	29F.1A12	
Applications:	FCM			
Reactivity:	Mouse			

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

PD-1; Programmed Death-1;		
Q02242		
CD279, also known as programmed death-1 (PD-1), is a 50-55 kD glycoprotein belonging to the CD28 family of the Ig superfamily. PD-1 is expressed on activated splenic T and B cells and thymocytes. It is induced on activated myeloid cells as well. PD-1 is involved in lymphocyte clonal selection and peripheral tolerance through binding its ligands, B7-H1 (PD-L1) and B7-DC (PD-L2). It has been reported that PD-1 and PD-L1 interactions are critical to positive selection and play a role in shaping the T cell repertoire. PD-L1 negative costimulation is essential for prolonged survival of intratesticular islet allografts.		
Liquid	APC Excitation and Emission Spectra	
APC	100 - 80 -	
25µg, 100µg		
Rat	60 (%)	
Rat IgG2a, κ	20 0 350 400 450 500 550 550 600 650 700 750 800 850 Wavelength (nm) Ex:650 nm	
	Q02242 CD279, also known as programm to the CD28 family of the Ig supe cells and thymocytes. It is induce lymphocyte clonal selection and (PD-L1) and B7-DC (PD-L2). It h critical to positive selection and pl costimulation is essential for proto Liquid APC 25µg, 100µg Rat	

Isotype Control: APC Rat IgG2a, κ Isotype Control[2A3] [Product AGEL1634]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- RecommendedEach lot of this antibody is quality control tested by flow cytometric analysis. Please check
your vial before the experiment. Since applications vary, the appropriate dilutions must be
determined for individual use. We suggest each investigator should titrate the reagent to
obtain optimal results [The recommended concentration is 0.1-1 μg/106 cells in 100 μL
volume].