

#### **Product Datasheet**

# **Low Endotoxin Purified Anti-Human** CD274/PD-L1 Antibody [29E.2A3]

Catalogue Code: AGEL1645

## Antibody Data

**Product SKU: AGEL1645** Clone: 29E.2A3

**Applications:** FCM;Block;FA

Reactivity: Human

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information:**

**Alternate Names:** B7-H1; PD-L1; Programmed cell death ligand 1;B7 homolog 1;B7-H; B7H1; PDL1;

PDCD1L1; PDCD1LG1;

**Uniprot ID:** Q9NZQ7

Background: CD274, also known as PD-L1 and B7-H1, is type I transmembrane glycoprotein that serves

as a ligand for CD279 (PD-1). This interaction is believed to regulate the balance between the stimulatory and inhibitory signals needed for responses to microbes and maintenance of self-tolerance. CD274 is involved in the costimulation of T cell proliferation and IL-10 and IFN-y production in an IL-2-dependent and CD279-independent manner. Conflicting data has shown that CD274 can inhibit T cell proliferation and cytokine production, and alternatively, enhance T cell activation. Other studies suggest that CD274 may signal bidirectionally, raising interesting implications for its expression in a wide variety of cell

types, including T and B cells, antigen-presenting cells, and nonhematopoietic cells.

Form: Liquid

Conjugation: None (AF/LE)

Size: 50µg, 500µg, 1mg

**Host Species:** Mouse

Isotype: Mouse IgG2b, κ

**Isotype Control:** AF/LE Purified Mouse IgG2b, K Isotype Control[MPC-11] [Product AGEL1645]

0.2 µm filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers Storage Buffer:

or stabilizers. Endotoxin level is < 2 EU/mg as Determined by LAL gel clotting assay.

Shipping: Biological ice pack at 4°C



**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to

light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial

contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 0.5~\mu g$  per 106 cells in 100  $\mu L$  volume or 100  $\mu L$  of whole blood. It is recommended that the reagent be titrated for optimal

performance for each application.