

Product Datasheet **PE Anti-Human CD34 Antibody [581]** Catalogue Code: AGEL1734

Antibody Data


Product SKU:	AGEL1734	Clone:	581
Applications:	FCM		
Reactivity:	Human		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	Gp105-120; My10;		
Uniprot ID:	P28906		
Background:	CD34, also known as gp105-120, is a type I monomeric sialomucin-like glycophosphoprotein with an approximate molecular weight of 105-120 kD. Selectively expressed on the majority of hematopoietic stem/progenitor cells, bone marrow stromal cells, capillary endothelial cells, embryonic fibroblasts, and some nervous tissue, CD34 is a commonly used marker to identify human hematopoietic stem/progenitor cells. According to the differential sensitivity to enzymatic cleavage, four groups of epitopes of CD34 have been described. CD34 mediates cell adhesion and lymphocytes homing through binding to L-selectin and E-selectin ligands.		
Form:	Liquid	PE Excitation and Emission Spectra	
Conjugation:	PE	80 -	
Size:	20 Tests, 100 Tests, 200 Tests		
Host Species:	Mouse	Weight and the second s	
Isotype:	Mouse IgG1, κ	20 0 350 400 450 500 550 600 650 700 Wavelength (nm)	



Isotype Control: PE Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL1734]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.