

Product Datasheet **PerCP Anti-Human CD11b Antibody [ICRF44]** Catalogue Code: AGEL1752

## Antibody Data


Product SKU:	AGEL1752	Clone:	ICRF44
Applications:	FCM		
Reactivity:	Human		

## Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

## Product Information:

Alternate Names: Uniprot ID: Background:	<ul> <li>Integrin alpha-M;Itgam;CD11 antigen-like family member B;CR-3 alpha chain;Leukocyte adhesion receptor MO1;CD11b; P11215</li> <li>CD11b is a 165-170 kD type I transmembrane glycoprotein also known as αMintegrin, Mac-1, CR3, and C3biR. CD11b non-covalently associates with integrin β2(CD18) and is expressed on granulocytes, monocytes/macrophages, dendritic cells, NK cells, and subsets of T and B cells. CD11b/CD18 is critical for the transendothelial migration of monocytes and neutrophils. It is also involved in granulocyte adhesion, phagocytosis, and neutrophil activation. CD11b/CD18 interacts with ICAM-1 (CD54), ICAM-2 (CD102), ICAM-4, CD14, CD23, heparin, iC3b, fibrinogen, and factor X.</li> </ul>		
Form:	Liquid	PerCP Excitation and Emission Spectra	
Conjugation:	PerCP	100	
Size:	20 Tests, 100 Tests, 200 Tests	80 - (2) (2) (3) (4) (4) (4) (4) (4) (4) (4) (4	
Host Species:	Mouse	40	
Isotype:	Mouse IgG1, κ	20 0 350 400 450 500 550 600 650 700 750 800 Wavelength (nm)	



**Isotype Control:** PerCP Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL1752]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.