

Product Datasheet **FITC Anti-Human CD61 Antibody [VI-PL2]** Catalogue Code: AGEL1825

## Antibody Data

Product SKU:	AGEL1825	Clone:	VI-PL2
Applications:	FCM		
Reactivity:	Human		

## Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

## **Product Information:**

Alternate Names:	GP3A;GPIIIa;ITGB3;Integrin beta-	3;
Uniprot ID:	P05106	
Background:	transmembrane glycoprotein. It is a gpIIb (CD41) to form CD41/CD6 (CD51/CD61) integrin. CD41/CD6 plays a role in platelet activation fibronectin, vWF, and other R expressed on platelets, osteocla	and glycoprotein IIIa (gpIIIa), is a 90 kD type I integral a member of the integrin family, associating with platelet I complex and with integrin $\alpha V$ (CD51) to form $\alpha V/\beta 3$ of is expressed on platelets and megakaryocytes, and and aggregation through interaction with fibrinogen, GD-containing adhesion molecules. CD51/CD61 is sts, fibroblasts, macrophages, and some tumor cells in adenovirus infection through binding to RGD motif in
Form:	المستنط	
i onn.	Liquid	FITC Excitation and Emission Spectra
Conjugation:	FITC	FITC Excitation and Emission Spectra
		100 -
Conjugation:	FITC	



**Isotype Control:** FITC Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL1825]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.