

#### **Product Datasheet**

# PE Anti-Human CD28 Antibody [CD28.2]

Catalogue Code: AGEL2021

## **Antibody Data**

Product SKU: AGEL2021 Clone: CD28.2

Applications: FCM

Reactivity: Human

### **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

## **Product Information:**

**Alternate Names:** PAS2;MNS;HGpMiX;GYPA;GPSAT;

Uniprot ID: P10747

Background: CD28 is a 44 kD disulfide-linked homodimeric type I glycoprotein. It is a member of the

immunoglobulin superfamily and is also known as T44 or Tp44. CD28 is expressed on most T lineage cells, NK cell subsets, and plasma cells. CD28 binds both CD80 and CD86 using a highly conserved motif MYPPY in the CDR3-like loop. CD28 is considered a major co-stimulatory molecule, inducing T lymphocyte activation and IL-2 synthesis, and preventing cell death. In vitro studies indicate that ligation of CD28 on T cells by CD80 and CD86 on antigen presenting cells provides a costimulatory signal required for T cell

activation and proliferation.


Form: Liquid

Conjugation: PE

Size: 20 Tests, 100 Tests, 200 Tests

Host Species: Mouse

**Isotype:** Mouse IgG1, κ



Ex:495;565 nm; Em:575 nm

**Isotype Control:** PE Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL2021]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C



Stability & Storage:

Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu L$  of antibody per test (million cells in 100  $\mu L$  staining volume or per 100  $\mu L$  of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.