

Low Endotoxin Purified Anti-Human/Mouse IL-5 Antibody [TRFK5]
 Catalogue Code: AGEL2062

Antibody Data

Product SKU:	AGEL2062	Clone:	TRFK5
Applications:	ICFCM;Depletion;Neut		
Reactivity:	Human;Mouse		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	Interleukin-5;IL-5;B-cell differentiation factor I;T-cell replacing factor;TRF;
Uniprot ID:	P05113 P04401
Background:	IL-5 is a homodimeric, disulphide-linked protein produced by T-cells. Monomeric human IL-5 is a 126 amino acid protein with a reported molecular weight of 26 kD for the homodimeric protein. Mouse and human IL-5 are approximately 70% identical. IL-5 has been shown to promote the growth of immature hematopoietic BFU-E progenitors, stimulate the activation and differentiation of eosinophils, and promote the generation of cytotoxic lymphocytes. Mouse IL-5 induces the differentiation and proliferation of pre-activated B-cells and stimulates the production and secretion of IgM and IgA by B-cells stimulated with bacterial endotoxin. The TRFK5 antibody can neutralize the bioactivity of natural or recombinant IL-5.
Form:	Liquid
Conjugation:	None (AF/LE)
Size:	50µg, 500µg, 1mg
Host Species:	Rat
Isotype:	Rat IgG1, κ
Isotype Control:	AF/LE Purified Rat IgG1, κ Isotype Control[HRPN] [Product AGEL2062]
Storage Buffer:	0.2 μm filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers or stabilizers. Endotoxin level is < 2 EU/mg as Determined by LAL gel clotting assay.
Shipping:	Biological ice pack at 4°C

Stability & Storage: Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

Recommended Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \mu\text{g}$ per 10^6 cells in 100 μL volume or 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
