

Product Datasheet **GenieFluor 647 Anti-Human CD44 Antibody [Hermes-1]** Catalogue Code: AGEL2107

## Antibody Data

Product SKU:	AGEL2107	Clone:	Hermes-1	
Applications:	FCM			
Reactivity:	Human			

## Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

## **Product Information:**

Alternate Names:	LHR;PGP-I;CDw44;Epican;HUTCH-I;			
Uniprot ID:	P16070			
Background:	CD44 is a 80-95 kD glycoprotein also known as Hermes, Pgp1, H-CAM, or HUTCH. It is expressed on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. As B and T cells become activated or progress to the memory stage, CD44 expression increases from a low or mid level of intensity to high expression levels. Thus, CD44 has been reported to be a valuable marker for memory cell subsets. CD44 is an adhesion molecule involved in leukocyte attachment to and rolling on endothelial cells, homing to peripheral lymphoid organs and to the sites of inflammation, and leukocyte aggregation.			
Form:	Liquid	647 Excitation and Emission Spectra		
Conjugation:	Genie Fluor647	100 -		
Size:	20 Tests, 100 Tests, 200 Tests	80 - E		
Host Species:	Rat	(%) po 100		
Isotype:	Rat IgG2a, к	$\frac{1}{9} \frac{40}{350} \frac{1}{400} \frac{1}{450} \frac{1}{500} \frac{1}{550} \frac{1}{600} \frac{1}{650} \frac{1}{700} \frac{1}{750} \frac{1}{800} \frac{1}{850} \frac{1}{850} \frac{1}{800} \frac{1}{$		

**Isotype Control:** Genie Fluor 647 Rat IgG2a, κ Isotype Control[2A3] [Product AGEL2107]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.