

#### **Product Datasheet**

# FITC Anti-Mouse CD45R/B220 Antibody [RA3.3A 1/6.1]

Catalogue Code: AGEL2214

### Antibody Data

Product SKU: AGEL2214 Clone: RA3.3A 1/6.1

Applications: FCM

Reactivity: Mouse

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information:**

Alternate Names: B220;

Uniprot ID: -

**Background**: CD45R, also known as B220, is an isoform of CD45. It is a member of the protein tyrosine

phosphatase (PTP) family with a molecular weight of approximately 180-240 kD. CD45R is expressed on B cells (at all developmental stages from pro-B cells through mature B cells), activated B cells, and subsets of T and NK cells. CD45R (B220) is also expressed on a subset of abnormal T cells involved in the pathogenesis of systemic autoimmunity in MRL-Fas<sup>|pr</sup> and MRL-Fas<sup>gld</sup> mice. It plays a critical role in TCR and BCR signaling. The primary ligands for CD45 are galectin-1, CD2, CD3, and CD4. CD45R is commonly used as a pan-B cell marker; however, CD19 may be more

appropriate for B cell specificity.


Form: Liquid

Conjugation: FITC

Size: 25µg, 100µg

Host Species: Rat

**Isotype:** Rat IgM, κ



Ex:490 nm; Em:530 nm

**Isotype Control:** FITC Rat IgM, κ Isotype Control[RTK2118] [Product AGEL2214]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C



**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to

light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial

contents. This product is guaranteed up to one year from purchase.

Recommended Usage:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu$ g/106 cells in 100  $\mu$ L volume].