

Product Datasheet **APC Anti-Mouse TCRβ Antibody [H57- 597 (HB218)]** Catalogue Code: AGEL2244

## Antibody Data

Product SKU:	AGEL2244	Clone:	H57-597 (HB218)
Applications:	FCM		
Reactivity:	Mouse		

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

## **Product Information:**

Alternate Names: Uniprot ID: Background:	TCR- $\beta$ chain; TCR- $\beta$ ; $\beta$ -TCR; - T cell receptor (TCR) is a heterodimer consisting of an $\alpha$ and a $\beta$ chain (TCR $\alpha/\beta$ ) or a $\gamma$ and a $\delta$ chain (TCR $\gamma/\delta$ ). TCR- $\beta$ is a member of the immunoglobulin superfamily and a component of the CD3/TCR complex (along with TCR- $\alpha$ ). It is expressed on $\alpha/\beta$ TCR- bearing T cells and thymocytes. The CD3/TCR complex plays a key role in antigen recognition, signal transduction, and T cell activation.		
Form:	Liquid	APC Excitation and Emission Spectra	
Conjugation:	APC	100 -	
Size:	50 Tests, 100 Tests, 200 Tests	80 - 30 - 10 -	
Host Species:	Armenian Hamster	A0 - 09 - 09 - 09 - 09 - 09 - 09 - 09 -	
Isotype:	Armenian Hamster IgG	20 0 350 400 450 500 550 550 600 650 700 750 800 850 Wavelength (nm) Ex:650 nm; Em:660 nm	
Isotype Control:	APC Armenian Hamster IaG Isotype Control[PIP] [Product AGEL2244]		

Isotype Control: APC Armenian Hamster IgG Isotype Control[PIP] [Product AGEL2244]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.