

Product Datasheet **PE/Cyanine5 Anti-Human CD161 Antibody [HP-3G10]** Catalogue Code: AGEL2280

Antibody Data

Product SKU:	AGEL2280	Clone:	HP-3G10	
Applications:	FCM			
Reactivity:	Human			

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	CLEC5B; NKRP1A;NKR-P1A;HNKR-P1a;KLRB1;		
Uniprot ID:	Q12918		
Background:	CD161 is a type II transmembrane glycoprotein, also known as NKR-P1A, that is expressed as a 40-44 kD homodimer. It is a member of the C-type lectin superfamily. CD161 is expressed on a majority of NK cells, NKT cells, and subsets of peripheral T cells and CD3+ thymocytes. It has been reported that Th17 cells are a subpopulation of CD4+CD161+CCR6+ cells. While the biological function of CD161 is not clear, it has been suggested to serve either as a stimulatory receptor or to inhibit NK cell-mediated cytotoxicity and cytokine production. LLT-1 (lectin-like transcript-1, also named as osteoclast inhibitory lectin or CLEC2D) is the ligand of CD161.		
Form:	Liquid	PE/Cyanine5 Excitation and Emission Spectra	
Conjugation:	PE/Cyanine 5	100	
Size:	20 Tests, 100 Tests, 200 Tests		
Host Species:	Mouse	S 00	
Isotype:	Mouse IgG1, κ	20 0 350 400 450 500 550 500 550 680 650 700 750 800 Wavelength (nm)	
		Ex:495;565;655 nm; Em:670 nm	
Isotype Control:	-		
Storage Buffer	Phoenbate buffered solution $nH 7.2$ containing 0.00% stabilizer and 1% protein protectant		

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.