

Product Datasheet **FITC Anti-Human CD39 Antibody [A1]** Catalogue Code: AGEL2288

Antibody Data

Product SKU:	AGEL2288	Clone:	A1
Applications:	FCM		
Reactivity:	Human		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	SPG64;NTPDase1;ATPDase;CD 39);	
Uniprot ID:	P49961		
Background:	Human CD39 is an integral membrane protein with two transmembrane domains. It exists as a homotetramer. Expression of CD39 is found on activated lymphocytes, a subset of T cells and B cells, and dendritic cells with weak staining on monocytes and granulocytes. CD39 and CD73 have been found on regulatory T cells, specifically the effector/memory like T cells. CD39 can hydrolyze both nucleoside triphosphates and diphosphates. CD39 is the dominant ecto nucleotidase of vascular and placental trophoblastic tissues and appears to modulate the functional expression of type 2 purinergic (P2) G protein coupled receptors (GPCRs). CD39 has intrinsic ecto-ATPase activity. Expression of CD39 is induced on T cells and increased on B cells as a late activation antigen. Product Details		
Form:	Liquid	FITC Excitation and Emission Spectra	
Conjugation:	FITC		
		80 - / / / /	
Size:	20 Tests, 100 Tests, 200 Tests	80 - (%) D 5 60 -	
Size: Host Species:	20 Tests, 100 Tests, 200 Tests Mouse	8 / /	



Isotype Control: FITC Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL2288]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.