

Product Datasheet **APC Anti-Human CD3 Antibody [UCHT1]** Catalogue Code: AGEL2466

## Antibody Data

| Product SKU:  | AGEL2466 | Clone: | UCHT1 |
|---------------|----------|--------|-------|
| Applications: | FCM      |        |       |
| Reactivity:   | Human    |        |       |

## **Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.

## Product Information:

| Alternate Names:<br>Uniprot ID:<br>Background: | T-cell surface glycoprotein CD3 epsilon chain;CD3E;T-cell surface antigen T3/Leu-4 epsilon chain;CD3e;CD3E;T3E;<br>P07766<br>CD3ɛ is a 20 kD chain of the CD3/T cell receptor (TCR) complex, which is composed of two CD3ɛ, one CD3γ, one CD3δ, one CD3ζ (CD247), and a T cell receptor ( $\alpha/\beta$ or $\gamma/\delta$ ) heterodimer. It is found on all mature T lymphocytes, NK T cells, and some thymocytes. CD3, also known as T3, is a member of the immunoglobulin superfamily that plays a role in antigen recognition, signal transduction, and T cell activation. |   |  |
|--|---|---|--|
| Form:  | Liquid  | APC Excitation and Emission Spectra                             |  |
| Conjugation:                                   | APC   | 80  |  |
| Size:  | 20 Tests, 100 Tests, 200 Tests  |   |  |
| Host Species:                                  | Mouse   | 60 - 50 - 50 - 50 - 50 - 50 - 50 - 50 -                         |  |
| Isotype:                                       | Mouse IgG1, κ   | 20<br>350<br>400<br>450<br>550<br>550<br>550<br>550<br>550<br>5 |  |

**Isotype Control:** APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL2466]

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.