

Product Datasheet **APC Anti-Human CD279/PD-1 Antibody** [EH12.2H7] Catalogue Code: AGEL2724

Antibody Data

Product SKU:	AGEL2724	Clone:	EH12.2H7
Applications:	FCM		
Reactivity:	Human		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names:	PDCD1;PD1;Protein PD-1;hPD-1;		
Uniprot ID:	Q15116		
Background:	Programmed cell death 1 (PD-1), also known as CD279, is a 55 kD member of the immunoglobulin superfamily. CD279 contains the immunoreceptor tyrosine-based inhibitory motif (ITIM) in the cytoplasmic region and plays a key role in peripheral tolerance and autoimmune disease. CD279 is expressed predominantly on activated T cells, B cells, and myeloid cells. PD-L1 and PD-L2 are ligands of CD279 (PD-1) and are members of the B7 gene family. Evidence suggests overlapping functions for these two PD-1 ligands and their constitutive expression on some normal tissues and upregulation on activated antigen-presenting cells. Interaction of CD279 ligands results in inhibition of T cell proliferation and cytokine secretion.		
Form:	Liquid	APC Excitation and Emission Spectra	
Conjugation:	APC	80 -	
Size:	20 Tests, 100 Tests, 200 Tests		
Host Species:	Mouse	Normalized (%)	
Isotype:	Mouse IgG1, κ	20	
		350 400 450 500 550 600 650 700 750 800 850 Wavelength (nm)	
		Ex:650 nm; Em:660 nm	
Isotype Control:	APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product AGEL2724]		

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.