

**Antibody Data**

<b>Product SKU:</b>	<b>AGEL2797</b>	<b>Clone:</b>	<b>3A8</b>
<b>Applications:</b>	<b>FCM</b>		
<b>Reactivity:</b>	<b>Human</b>		

**Important Note:**

Centrifuge before opening to ensure complete recovery of vial contents.


**Product Information:**

**Alternate Names:** Tumor necrosis factor receptor superfamily member 5;Cd40;B-cell surface antigen CD40;Bp50;CD40L receptor;CD40;Tnfrsf5; P25942

**Uniprot ID:**

**Background:** CD40 is a 48 kD type I glycoprotein also known as BP50. It is a member of the TNFR superfamily primarily expressed on B cells, macrophages, follicular dendritic cells, endothelial cells, fibroblasts, and at low levels on plasma cells. CD40 has been reported to be involved in B cell differentiation, costimulation, isotype class-switching, and protection of B cells from apoptosis. Additionally, CD40 is important for T cell-B cell interactions. The ligand of CD40 is CD154 (CD40 ligand). The 5C3 antibody has been reported to promote B cell proliferation in the presence of anti-IgM, IL-4 or PMA, partially blocking CD40 binding to CD40L, and B cells rescue from apoptosis.

<b>Form:</b>	Liquid
<b>Conjugation:</b>	Genie Fluor488
<b>Size:</b>	20 Tests, 100 Tests, 200 Tests
<b>Host Species:</b>	Mouse
<b>Isotype:</b>	Mouse IgG2a, κ



<b>Isotype Control:</b>	Genie Fluor 488 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product AGEL2797]
<b>Storage Buffer:</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
<b>Shipping:</b>	Biological ice pack at 4°C

**Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.

**Recommended Usage:** Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.