

Product Datasheet **FITC Anti-Human CD44 Antibody [P2A1]** Catalogue Code: AGEL2798

Antibody Data

Applications: FCM	Product SKU:	AGEL2798	Clone:	P2A1
	Applications:	FCM		
Reactivity: Human	Reactivity:	Human		

Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

Product Information:

Alternate Names: Uniprot ID:	CD44 antigen;CD44;CDw44;Epica glycoprotein I;PGP-I;CD44;LHR; MD P16070		
Background:	CD44 is a 80-95 kD glycoprotein also known as Hermes, Pgp1, H-CAM, or HUTCH. It is expressed on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. As B and T cells become activated or progress to the memory stage, CD44 expression increases from a low or mid level of intensity to high expression levels. Thus, CD44 has been reported to be a valuable marker for memory cell subsets. CD44 is an adhesion molecule involved in leukocyte attachment to and rolling on endothelial cells, homing to peripheral lymphoid organs and to the sites of inflammation, and leukocyte aggregation.		
Form:	Liquid	FITC Excitation and Emission Spectra	
Conjugation:	FITC		
Size:	20 Tests, 100 Tests, 200 Tests	80 - S = 0 -	
Host Species: N	Mouse	\$ 60 - 11 TITE 40 -	
Isotype:	Mouse IgG2a, к	2 0 0 350 400 450 500 550 600 650 700 Wavelength (nm)	

Isotype Control: FITC Mouse IgG2a, κ Isotype Control[C1.18.4] [Product AGEL2798]

Storage Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Ex:490 nm; Em:530 nm

Shipping: Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.