

Product Datasheet **GenieFluor Red 780 Anti-Human CD8a Antibody [OKT-8]** Catalogue Code: AGEL2979

## Antibody Data

Product SKU:	AGEL2979	Clone:	OKT-8
Applications:	FCM		
Reactivity:	Human		

## Important Note:

Centrifuge before opening to ensure complete recovery of vial contents.

## Product Information:

Alternate Names: Uniprot ID:	T-cell surface glycoprotein CD8 alpha chain;CD8A;T-lymphocyte differentiation antigen T8/Leu-2;MAL; P01732		
Background:	CD8a is a 32-34 kD type I glycoprotein. It forms a homodimer (CD8a/a) or heterodimer (CD8a/b) with CD8b. CD8, also known as T8 and Leu2, is a member of the immunoglobulin superfamily found on the majority of thymocytes, a subset of peripheral blood T cells, and NK cells (which express almost exclusively CD8a homodimers). CD8 acts as a co-receptor with MHC class I-restricted T cell receptors in antigen recognition and T cell activation and has been shown to play a role in thymic differentiation. Two domains in CD8a are important for function: the extracellular IgSF domain binds the $\alpha$ 3 domain of MHC class I and the cytoplasmic CXCP motif binds the tyrosine kinase p56 Lck.		
Form:	Liquid	'80 Excitation and Emission Spectra	
Conjugation:	Genie FluorRed 780	100	
Size:	20 Tests, 100 Tests, 200 Tests	80 -	
Host Species:	Mouse	(%) 60 . 10 . 40 .	
Isotype:	Mouse IgG2a, к	E 40 20 	
Isotype Control:	Genie Fluor Red 780 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product AGEL2979]		
Storage Buffer	Phosphate buffered solution pH 7.2 containing 0.00% stabilizer and 1% protein protectant		

**Storage Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping:** Biological ice pack at 4°C



- **Stability & Storage:** Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. Centrifuge before opening to ensure complete recovery of vial contents. This product is guaranteed up to one year from purchase.
- **Recommended** Usage: Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.